



CRINONE is administered vaginally, once or twice daily. Each applicator is individually packaged, for one-time use. After administration, a small amount of CRINONE will be left in the tube—you will still be receiving the proper dose. Once in the vagina, CRINONE coats the vaginal wall, where the progesterone is absorbed and transported directly to the uterus.

Instructions for using CRINONE

- Remove the applicator from the sealed wrapper. DO NOT remove the twist-off tab at this time
- Hold the applicator between the thumb and forefinger along the seam on the thick end
- Shake down 3 to 4 times (like a thermometer) to ensure that the contents are at the thin end of the applicator



- Hold the applicator by the flat section of the thick end
- Twist off and throw away the tab at the thin end. Remember to twist off completely—do not pull the tab off
- DO NOT squeeze the thick end while twisting the tab. This could force some gel to be released before it is inserted into the vagina



- The applicator may be inserted into your vagina while you are in a sitting position or when lying on your back with your knees bent
- Gently insert the thin end well into your vagina



- Squeeze the thick end of the applicator to deposit CRINONE into the vagina
- Remove the applicator and throw it away in a waste container



Disclaimer:

Instruction for using Crinone information is provided by http://www.crinoneusa.com/patient/how_to_use_crinone.asp and http://www.emdserono.ca/cmgeemdserono.ca/en/images/CRINONE_tcm463_84822.pdf?Version= (page 12, 13) for reference purpose. Please consult your Fertility physician, nurses or pharmacists for your dosage information and professional instructions and training for using Crinone.